

PAPER • OPEN ACCESS

Psychochemical and Organoleptic Characteristics of Colostrum Kefir as Antibacterial

Recent citations

- [Antibacterial activity of colostrum kefir against foodborne pathogen bacteria](#)
N Windayani *et al*

To cite this article: N Windayani *et al* 2019 *J. Phys.: Conf. Ser.* **1175** 012016

View the [article online](#) for updates and enhancements.

The advertisement banner features a collage of colorful book covers on the left, including titles like 'Infrared Imaging' and 'Physics and Quantum Physics'. On the right, the text reads: 'IOP | ebooks™ Bringing together innovative digital publishing with leading authors from the global scientific community. Start exploring the collection—download the first chapter of every title for free.'

IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Psychochemical and Organoleptic Characteristics of Colostrum Kefir as Antibacterial

N Windayani^{1*}, T Kurniati², and M Listiawati²

¹Department of Chemistry Education, ²Department of Biology Education
Faculty of Education and Teacher Training, UIN Sunan Gunung Djati Bandung,
Bandung, Indonesia

*nenengwinda.ftk@uinsgd.ac.id

Abstract. This study aims to create and analyze psychochemical, and organoleptic characteristics of colostrum kefir as an antibacterial. Colostrum kefir is made using a three-factor completely randomized design method with three replications. Psychochemical characterization includes pH, specific gravity, lactic acid content, and alcohol content, while the aspects of organoleptic testing by panelist include color, aroma, taste, texture, and viscosity. This study has succeeded in making colostrum kefir under the optimum condition, at 20% grain kefir, with 48-hour fermentation at 28 °C with the characteristics as follows; pH 3,31-3,65; lactic acid content 1,96-2,83 %; alcohol content 0,38-0,59 %; and Specific gravity 1,030-1,046 mg/mL. Moreover, the results of organoleptic testing show that the colostrum kefir obtained is yellowish orange, sour, rather soft and thick and has a rather typical kefir aroma.

1. Introduction

Recently, there has been a trend toward the increase in demand for functional food globally, along with the rise in awareness to reduce the risk of disease and improve the quality of health (1). Demand for functional food in Europe was reported to reach 14%, the United States 38%, while Japan as first ranks by 40% (2). Indonesia itself listed as one of the most significant functional food markets in Southeast Asia (1). Several types of functional food have been distributed to the community, including dairy products, probiotics, yogurt, kefir, and koumiss (2). Milk kefir has long been believed to have various health benefits (3) from its antioxidant (4), anticancer (5), antidiabetic (6), antihypertensive (7), antimicrobial (8), and anti-odor (for petai) (9) activity to its skin care properties (10). Nowadays, there is even increasing attention to milk kefir, including its production at industrial scale (11).

The types of ingredients used as milk kefir include cow milk, soy milk (12), goat milk (13), and camel milk (14). However, there has been no report to date about colostrum kefir namely colostrum milk fermented with kefir probiotics. Whereas, colostrum has many features including very high protein content (15) mainly composed of immunoglobulin (16), lactoferrin (17), casein, growth factors, enzymes, cytokines, nucleosides, nucleotides, vitamins, minerals, and lipids (15). The benefits of colostrum for health include its ability to modulate the body immune system (18), and its antiviral (19) and antimicrobial activities (20).

However, colostrum is very susceptible to microbial contamination (21) and also not resistant to heating due to its high protein content (15). Therefore proper treatment is required so that its function can be utilized optimally for human health. One of the efforts possible to make is to ferment colostrum using kefir probiotics. Hence, this study studies further regarding the optimum fermentation condition



and psychochemical and organoleptic characteristics of colostrum kefir as the materials for testing its antibacterial properties.

2. Methods

2.1. Colostrum material collection

The raw material of colostrum obtained from cattle farms around the foot of Manglayang Mountain, namely the first milk colostrum (P1), 24 hours after postpartum.

2.2. Kefir grain collection

Kefir grain obtained through purchases from activists of Indonesian Kefir Community in Bandung. This kefir grain has been previously identified in the UNPAD Microbiology Laboratory, Jatiningor, to determine the type and number of microbes composing the grain.

2.3. The making of colostrum kefir using kefir grain as the starter

The making of kefir colostrum was done by modifying the previous work (22). Briefly, 250 mL of P1 colostrum is fermented by inoculating kefir grain 10%, 15%, and 20%, respectively in food grade plastic fermenters then stirred slowly using a stainless stirrer until homogeneous. Next, it is incubated for 24, 48, and 72 hours, with occasionally stirring after 12, 24, and 48 hours. The optimum fermentation condition is determined using three-factor completely randomized design with three replications based on kefir grain concentration and fermentation time. The first factor is the starter dose of kefir grain, namely 10%, 15%, and 20% and the second factor is the fermentation time, namely 24, 48, and 72 hours. The result of this stage is in the form of data of the optimum time and concentration of kefir grain to produce colostrum kefir.

2.4. Determination of the characteristics of fermented colostrum kefir

The psychochemical characteristics of colostrum kefir refer to the previous work (22), including pH, lactic acid content, alcohol content, and specific gravity. The organoleptic testing uses parameters including taste, aroma, color, texture, and viscosity (23).

3. Results and Discussion

3.1. The degree of acidity (pH)

The degree of acidity indicates the level of acidity (pH) of a product. The observations show that during the colostrum fermentation process there has been a change in pH as a sign of the fermentation process (7). The initial pH of Cow colostrum as a raw material in this fermentation is 5.25, and then after fermentation with the starter, i.e., kefir grain, 10%, 15%, and 20%, it decreases to 3.31 and 3.65. The formation of Lactic acid causes the acidity of kefir as the product of milk/lactose sugar decomposition (7) by lactic acid bacteria contained in kefir grains. The result of the average pH of colostrum kefir produced from each treatment presented in Table 1.

3.2. Lactic acid content

As previously stated, lactic acid contained in kefir is produced from the decomposition of lactose by lactic acid bacteria. This also causes milk kefir in general and colostrum kefir in this work to be acidic indicated by pH ranging from 3.35 to 3.65. Furthermore, the Lactic content is calculated based on the titration results of each sample using NaOH 1N. In this study, the highest content of Lactic acid is produced by colostrum kefir sample F, i.e. 2.83% while sample A i.e. 1.96% produces the lowest content. Meanwhile, the quality requirement of kefir according to CODEX-STAN 243-2003 is at least 0.6% (24), meaning that the colostrum kefir obtained has met the criterion.

Table 1. Psychochemical characteristics of colostrum kefir

Sample ^a	Parameter			
	pH	Lactic acid(%)	Alcohol (%)	Specific gravity (g/mL)
A	3,65	1,96	0,39	1,030
B	3,62	1,97	0,38	1,033
C	3,61	1,98	0,38	1,038
D	3,35	2,64	0,50	1,041
E	3,40	2,50	0,55	1,046
F	3,31	2,83	0,46	1,040
G	3,40	2,52	0,57	1,043
H	3,41	2,50	0,52	1,043
I	3,40	2,46	0,57	1,042

^a A=d1w1, B= d2w1, C= d3w1, D= d1w2, E= d2w2, F= d3w2, G= d1w3, H= d2w3, I= d3w3

3.3. Alcohol content

Alcohol contained in kefir is produced from the decomposition of glucose by the *Saccharomyces cereviceae* yeast contained in kefir grain. Based on Table 1, in this study, the highest alcohol content is produced by colostrum kefir samples G and I, i.e., 0.57% while samples B and C provides the lowest content, i.e., 0.38%. It means that the amount of yeast in the sample with 20% concentration of kefir grain is more than the other two concentrations. According to CODEX STAN 243-2003, kefir contains at least 10⁴ CFU/g (24). The existence of Alcohol in kefir is a typical sign, distinguishing it from other fermented dairy products such as yogurt.

3.4. Specific gravity

Specific gravity is one of the physical parameters in milk fermentation. Based on Table 1, the specific gravity of kefir colostrum resulted in ranges from 1,030-1,046 g / mL. The colostrum kefir sample E shows the highest specific gravity, i.e., 1,046 g / mL while the lowest Specific gravity is demonstrated by sample A, i.e., 1,030 g / mL. Compared with the specific gravity of colostrum material used, i.e. 1,029 g / mL, the colostrum kefir resulted after fermentation in this study has a higher specific gravity than before fermentation. The data in Table 1 show that the lower the pH, the higher the specific gravity of the colostrum kefir produced. The previous report (25) states that the psychochemical characteristic of kefir products was determined by the condition of the growth of the kefir starter.

3.5. The organoleptic testing results of colostrum kefir

The organoleptic testing results of colostrum kefir done by seven panelists presented in Table 2. Based on the results of the testing, the panelists generally state that all colostrum kefir samples are yellowish orange, sour, slightly soft and thick and have a somewhat typical kefir aroma.

Table 2. Organoleptic testing of colostrum kefir

Assessment criteria	Sample									
	A	B	C	D	E	F	G	H	I	
Color	Very orange	0	2	0	0	0	0	0	0	0
	Yellowish orange	4	5	4	5	6	5	4	5	4
	A little orange	1	0	2	2	1	2	2	1	1
	Orange	2	0	1	0	0	0	1	1	2
	Yellow	0	0	0	0	0	0	0	0	0
Aroma	Very typical of kefir	1	2	0	0	0	0	1	1	0
	typical of kefir	0	2	2	0	1	1	1	1	1
	rather typical of kefir	5	3	3	6	4	5	5	4	6
	Typical of Milk	1	2	1	1	2	1	0	1	0

	Rather typical of butter	0	0	1	0	0	0	0	0	0
	Typical of butter	0	0	0	0	0	0	0	0	0
Taste	Very sour	0	0	0	0	0	0	0	1	1
	A little sour	1	2	2	1	2	1	1	1	1
	sour	5	4	3	5	4	6	5	4	5
	rather sour	1	1	2	1	1	0	1	1	0
	Not sour	0	0	0	0	0	0	0	0	0
	Not very sour	0	0	0	0	0	0	0	0	0
Texture	Very soft	0	0	0	0	0	0	0	0	0
	soft	3	0	1	0	1	1	0	0	0
	Rather soft	3	6	5	4	4	5	5	6	6
	Medium	1	0	1	2	1	1	2	1	0
	Rather hard	0	1	0	1	2	0	0	0	1
	hard	0	0	0	0	0	0	0	0	0
Viscosity	Very hard	0	0	0	0	0	0	0	0	0
	Very thick	0	0	3	1	1	2	1	1	2
	Thick	1	4	4	4	5	4	5	6	5
	Rather thick	2	3	0	2	1	1	0	1	0
	Medium	3	0	0	0	0	0	1	0	0
	A little thin	0	0	0	0	0	0	0	0	0
thin	1	0	0	0	0	0	0	0	0	
	Very thin	0	0	0	0	0	0	0	0	0

Compared with Indonesian national standards (SNI) for similar fermentation products, namely yogurt (SNI 2981: 2009), the organoleptic characteristics of colostrum kefir produced indicates criteria that meet the standard. Considering that there is no SNI for kefir and its derivatives to date, this study uses SNI for similar products namely yogurt. The comparison of colostrum kefir obtained in this study with the criteria for quality requirements of SNI 2981: 2009 presented in Table 3.

Table 3. Comparison of characteristics of colostrum kefir with SNI 2981:2009

Assessment criteria	colostrum kefir	SNI 2981:2009
Appearance	thick	Viscous liquid - Solid
odor / aroma	Rather typical of kefir	normal/typical
Taste	sour/typical	sour/typical
Consistency	homogeneous	homogeneous
Lactic acid content (%)	1,96-2,83	0,50-2,00

Furthermore, based on the results of this study, the bioactivity of the colostrum kefir obtained is tested as an antimicrobial, both for bacteria and for pathogenic fungi.

4. Conclusion

This study has succeeded in making colostrum kefir meeting the criteria for CODEX STAN 243-2003 and SNI 2981:2009 with the characteristics as follows: pH 3,31-3,65; Lactic acid content 1,96-2,83 %; Alcohol content 0,38-0,59 %; Specific gravity 1,030-1,046 mg/mL; and organoleptic attributes as follows: yellowish orange, sour, slightly soft and thick with a rather typical kefir aroma.

Acknowledgment

The authors would like to express a great appreciation and thanks to the Center for Research and Publishing (Puslitpen) UIN Sunan Gunung Djati Bandung for the 2018 Competitive Interdisciplinary Research Grants with Contract Number B-133/A3-29/Un.05/V.2/PP.00.9/03/2018, and Dapur Kefir Bandung for providing of cow colostrum and kefir grain.

References

1. Basu SK, Thomas JE, Acharya SN. Prospects for growth in global nutraceutical and functional food markets : A Canadian perspective. *Aust J Basic Appl Sci.* 2007;1(4):637–49.
2. Kozonov J, Povarova N. Leading Trends In The European Functional Food Market. *Холодильна Техніка Та Технологія* [Internet]. 2015;51(4):2–6. Available from: <http://journals.uran.ua/reftech/article/view/44785>
3. Gaware V, Kotade K, Dolas R, Dhamak K, Somwanshi S, Nikam V, et al. The magic of kefir: a review. *Pharmacologyonline.* 2011;1:376–86.
4. Chen Z, Shi J, Yang X, Nan B, Liu Y, Wang Z. Chemical and physical characteristics and antioxidant activities of the exopolysaccharide produced by Tibetan kefir grains during milk fermentation. *Int Dairy J.* 2015;43.
5. dos Reis SA, da Conceição LL, Siqueira NP, Rosa DD, da Silva LL, Peluzio M do CG. Review of the mechanisms of probiotic actions in the prevention of colorectal cancer. *Nutr Res.* 2017;37.
6. Judiono, Hadisaputro S, Ks I, Cahyono B, Suzery M. Effects of Clear Kefir on Biomolecular Aspects of Glycemic Status of Type 2 Diabetes Mellitus (T2DM) Patients in Bandung , West Java [Study on Human Blood Glucose , c Peptide and Insulin]. *Funct Foods Heal Dis.* 2014;4(8):340–8.
7. Febrisiantosa A, Priyo Purwanto B, Isnafia Arief I, Widyastuti Y. Karakteristik Fisik, Kimia, Mikrobiologi Whey Kefir Dan Aktivasnya Terhadap Penghambatan Angiotensin Converting Enzyme (Ace). *J Teknol dan Ind Pangan* [Internet]. 2013;24(2):147–53. Available from: <http://journal.ipb.ac.id/index.php/jtip/article/view/7694>
8. Al-Kalifawi EJ. Anti-Bacterial Activity of the Prophet Mohammad (SWS) Drink â€^{TM} s against Pathogenic Bacteria. *J Nat Sci Res.* 2015;5(4):90–9.
9. Kurniati T, Windayani N, Listiawati M. Anti-odor activity of milk kefir on organosulphur polysulfide cyclic compounds in petai (*parkia speciosa* hassk). *J Phys Conf Ser.* 2018;1013(1).
10. Chen MJ, Liu JR, Sheu JF, Lin CW, Chuang CL. Study on skin care properties of milk kefir whey. *Asian-Australasian J Anim Sci.* 2006;19(6):905–8.
11. M-Guedes KT, Magalhães KT, Schwan RF. Chemical and Therapeutic Aspects of Kefir. *Int J Pharmacovigil.* 2016;1(1):8–10.
12. Kesenkas Harun Nayil Dinkci, Kemal Seckin, Özer Kinik SG. Antioxidant Properties of Kefir Produced from Different Cow and Soy Milk Mixtures. *J Agric Sci.* 2011;17:253–9.
13. Nurliyani N, Sadewa A, Sunarti S. Kefir Properties Prepared with Goat Milk and Black Rice (*Oryza sativa* L.) Extract and its Influence on the Improvement of Pancreatic beta-Cells in Diabetic Rats. *Emirates J Food Agric* [Internet]. 2015;27(10):1. Available from: <http://www.ejfa.me/?mno=186533>
14. Kavas G. Kefirs Manufactured from Camel (*Camelus dromedarius*) Milk and Cow Milk : Comparison of some chemical and microbial properties. *Ital J Food Sci.* 2015;27:357–66.
15. McGrath BA, Fox PF, McSweeney PLH, Kelly AL. Composition and properties of bovine colostrum : a review. *Dairy Sci Technol.* 2016;96:133–58.
16. Hurley WL, Theil PK. Perspectives on immunoglobulins in colostrum and milk. *Nutrients.* 2011;3(4):442–74.
17. Ayar A, Sıçramaz H, Çetin İ. The Effect of Bovine Colostrum on the Lactic Flora of Yogurt and Kefir. *JSM Biotechnol Bioeng.* 2016;3(5):3–8.
18. Jørgensen ALW, Juul-Madsen HR, Stagsted J. Colostrum and bioactive, colostrum peptides differentially modulate the innate immune response of intestinal epithelial cells. *J Pept Sci.* 2010;16(1):21–30.
19. Waarts BL, Aneke OJC, Smit JM, Kimata K, Bittman R, Meijer DKF, et al. Antiviral activity of human lactoferrin: Inhibition of alphavirus interaction with heparan sulfate. *Virology.* 2005;333(2):284–92.
20. Berlov MN, Korableva ES, Andreeva Y V., Ovchinnikova T V., Kokryakov VN. Lactoferrin from canine neutrophils: Isolation and physicochemical and antimicrobial properties. *Biochem* [Internet]. 2007;72(4):445–51. Available from: <http://link.springer.com/10.1134/S0006297907040128>

21. Santos G, Thaís J, Hermelina F. *Revista Brasileira de Zootecnia* Nutritional and microbiological quality of bovine colostrum samples in Brazil. 2017;
22. Kurniati T, Windayani N, Listiawati M. Total Asam Laktat, Protein, Lemak, Karbohidrat, dan Serat Whey Kefir Susu Sapi Berdasarkan Konsentrasi Starter dan Waktu Fermentasi. In: *Seminar Nasional Bioteknologi UGM*. 2016. p. 1–11.
23. Surayah A, Sugiarto. Uji kimiawi dan organoleptik sebagai uji mutu yoghurt. *Pros Temu Tek Nas Tenaga Fungsional Pertan*. 2005;108–13.
24. World Health Organization and Food and Agriculture Organization of the United States. *Codex Standard for Fermented Milks - CODEX STAN-243-2003*. Codex Aliment [Internet]. 2011;56–8. Available from: <http://www.fao.org/docrep/015/i2085e/i2085e00.pdf>
25. Ismaiel AA, Ghaly MF, El-Naggar AK. Some physicochemical analyses of kefir produced under different fermentation conditions. *J Sci Ind Res (India)*. 2011;70(5):365–72.