

PN9

by Mohamad Agus Salim

Submission date: 21-Apr-2023 09:57AM (UTC+0700)

Submission ID: 2070923893

File name: Diseminasi_SAINTEK_2018_lengkap.pdf (1.96M)

Word count: 320

Character count: 2225



uin

Sertifikat

Nomor : B-122b/Un.05/III.7/PP.00.9/1/2018

diberikan Kepada :

Dr. Mohammad Agus Salim, Drs., M.P.

sebagai :


PEMATERI

dalam kegiatan Diseminasi Hasil Penelitian
Dosen Fakultas Sains dan Teknologi UIN Sunan Gunung Djati Bandung
pada tanggal, 23 Januari 2018

Dekan
Fakultas Sains dan Teknologi


Dr. H. Opik Taupik Kurahman
NIP. 19681214 199603 1 001

Ketua Pelaksana,


Adam Faroqi, MT.
NIP. 19740516 200912 1 001

Evaluation of Anticataract Potential of *Porphyridium cruentum* Biomass in Mice Lens

Diseminasi Hasil Penelitian Saintek

Bandung, 23 Januari 2018



uin
UNIVERSITAS ISLAM NEGERI
SUNAN GUNUNG DJATI
BANDUNG


Mohamad Agus Salim

Indonesia:


62 % water

**+/- 18.110
islands,
108.900 km
coastline**

**5100 km from
Aceh to
Papua**



Microalgae Culture



A microalgae from the family of red algae, rich in antioxidant compounds such as flavonoid, β carotene, vit E and C, phenolic compounds, chlorophyll and especially phycoerithrin.





Porphyridium cruentum



Cataract development is a gradual process but it can occur rapidly.




The remedy of cataract is surgery.
Problems of surgery: long waiting lists, costs, risks of complication and lack of technical equipment.



Backlog in Indonesia:

- cataract patients: 210,000 people/year
- only 80,000 people/year undergo surgery



Cataract risk factors



- age > 40 year



- genetic



- diabetes



- drugs



- sunlight



- smoking



- alcoholic beverages




It is estimated that a 10-year delay would reduce the need for surgery by 50%.






Research Objectives

- Study the potentiality of *P. cruentum* to protect against cataract formation in mice lens



- Develop an alternative method to surgery for the treatment of cataract

Contribution



Experimental Design

G1 : Aquabidest 2cc p.o)

G2 : Naphthalene 1 g/kg BW/day (0,5 g/kg BW/day in 3 days early)




G3 : P. cruentum 1g/kg BW/day

G4 : P. cruentum 1g/kg BW/day + Naphthalene 1g/kg BW/day



DEGREE OF OPACIFICATION

THE LENS SOLUBLE PROTEIN & WATER CONTENT



1

STATISTICAL ANALYSIS



Values are presented as mean \pm SEM. Results were compared by one-way ANOVA followed by Duncan's test. A value of $p < 0.001$ was considered significant

RESULTS


Group	Treatment	Soluble Protein (x 10 ⁻³ µg/ml)	Water Content (%)
G1	Control	0,48 ± 0,04	59 ± 4.67
G2	<i>P. cruentum</i>	0,49 ± 0,03	60 ± 7,31
G3	Naphthalene	0,39 ± 0,01	30 ± 10,54
G4	<i>P. cruentum</i> + Naphthalene	0,42 ± 0,02	53 ± 11,30

Group	Treatment	Degree of opacification
-------	-----------	-------------------------

G 1 :	Control	0
G 2 :	P. cruentum	0
G 3 :	Naphthalene	+++
G 4 :	P. cruentum +Naph	+




G1



G2



G3



G4

Conclusion

P. cruentum biomass can prevent the cataract progression in naphthalene-induced cataract models.

**Thank you for listening
I hope you found it interesting**

ORIGINALITY REPORT

12%
SIMILARITY INDEX

12%
INTERNET SOURCES

10%
PUBLICATIONS

0%
STUDENT PAPERS

PRIMARY SOURCES

1 www.ijpp.com **9%**
Internet Source

2 patents.google.com **3%**
Internet Source

Exclude quotes Off
Exclude bibliography On

Exclude matches Off